

blood

2012 119: 857-860
Prepublished online November 30, 2011;
doi:10.1182/blood-2011-03-341776

Regulation of hepcidin expression at high altitude

Nick P. Talbot, Samira Lakhal, Thomas G. Smith, Catherine Privat, Annabel H. Nickol, Maria Rivera-Ch, Fabiola León-Velarde, Keith L. Dorrington, David R. Mole and Peter A. Robbins

Updated information and services can be found at:
<http://bloodjournal.hematologylibrary.org/content/119/3/857.full.html>

Articles on similar topics can be found in the following Blood collections
[Brief Reports](#) (1509 articles)
[Red Cells, Iron, and Erythropoiesis](#) (326 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://bloodjournal.hematologylibrary.org/site/subscriptions/index.xhtml>



Brief report

Regulation of hepcidin expression at high altitude

*Nick P. Talbot,^{1,2} *Samira Lakhal,^{1,3} *Thomas G. Smith,^{1,4} Catherine Privat,⁵ Annabel H. Nickol,^{1,2} Maria Rivera-Ch,⁵ Fabiola León-Velarde,⁵ Keith L. Dorrington,^{1,4} David R. Mole,³ and Peter A. Robbins¹

¹Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom; ²Oxford Centre for Respiratory Medicine, Churchill Hospital, Oxford, United Kingdom; ³Henry Wellcome Building for Molecular Physiology, University of Oxford, Oxford, United Kingdom; ⁴Nuffield Department of Anaesthetics, John Radcliffe Hospital, Oxford, United Kingdom; and ⁵Laboratory of High Altitude Adaptation, Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Peru

Enhanced erythropoietic drive and iron deficiency both influence iron homeostasis through the suppression of the iron regulatory hormone hepcidin. Hypoxia also suppresses hepcidin through a mechanism that is unknown. We measured iron indices and plasma hepcidin levels in healthy volunteers during a 7-day

sojourn to high altitude (4340 m above sea level), with and without prior intravenous iron loading. Without prior iron loading, a rapid reduction in plasma hepcidin was observed that was almost complete by the second day at altitude. This occurred before any index of iron availability had changed. Prior iron loading de-

layed the decrease in hepcidin until after the transferrin saturation, but not the ferritin concentration, had normalized. We conclude that hepcidin suppression by the hypoxia of high altitude is not driven by a reduction in iron stores. (*Blood*. 2012;119(3):857-860)

Introduction

Dietary iron absorption was shown in the 1950s to be under the dual regulation of body iron stores and the prevailing rate of erythropoiesis, giving rise to the concept of the “store regulator” and the “erythroid regulator” in iron homeostasis.^{1,2} The downstream mediator of both regulators was later shown to be the iron-regulatory hormone hepcidin,³ which opposes iron absorption and recycling by binding and internalizing the cellular iron exporter ferroportin.⁴

The molecular mechanisms of hepcidin regulation are not completely understood. Oral iron loading stimulates hepcidin production in healthy humans.^{5,6} This may result in part from the binding of serum holotransferrin to transferrin receptor 1 in the liver, releasing the non-classical MHC class I molecule hereditary hemochromatosis protein (HFE) to interact with a holotransferrin-transferrin receptor 2 (TfR2) complex that up-regulates hepcidin expression.⁷⁻⁹ There is also evidence of a role for the membrane-bound protein hemojuvelin in the response to iron loading,^{5,10} and mutations in HFE, TfR2, or hemojuvelin can all cause hemochromatosis via inadequate hepcidin production.¹¹ The signal linking erythropoiesis to hepcidin suppression remains unknown. Candidates include erythropoietin (EPO) itself, soluble transferrin receptor (sTfR), and growth differentiation factor-15 (GDF15), which is released by maturing erythroblasts and reduces hepcidin mRNA expression in cultured hepatocytes.¹² Excessive serum GDF15 appears to underlie the paradoxical iron overload seen in patients with β -thalassemia.¹²

It is now known that hepcidin expression is also suppressed by hypoxia, but the mechanism is controversial. In mice, hypoxia-inducible factor was reported to regulate hepcidin expression via direct transcriptional suppression,¹³ but this finding has not been replicated in isolated hepatocytes.¹⁴ Hypoxia-inducible factor may,

however, contribute to hepcidin suppression indirectly via effects on the breakdown of hemojuvelin.^{15,16}

In vivo, hypoxia could also suppress hepcidin indirectly through erythropoiesis and enhanced iron use. Recently, a decrease in hepcidin was reported in healthy volunteers ascending gradually to high altitude.¹⁷ In this study, the authors observed a strong correlation between serum ferritin and hepcidin at each time point, and from this observation concluded that iron itself, or the kinetics of iron use in response to hypoxia, may be a signal for hepcidin down-regulation.

In the present study, we tested the hypothesis that the suppression of hepcidin by hypoxia arises through a reduction in iron availability. First, the changes in hepcidin and iron status on exposure to high altitude were examined at a greater temporal resolution than had been achieved in the previous report.¹⁷ Volunteers were transported from sea level to high altitude within an 8-hour period, and blood samples were taken daily during the first 3 days at high altitude. Secondly, a further group of volunteers underwent iron loading before travel to high altitude, to determine whether this would abrogate the reduction in hepcidin with hypoxia.

Methods

This study conformed to the principles of the Declaration of Helsinki and was approved by the Oxford Tropical Research Ethics Committee (Oxford, United Kingdom) and the Universidad Peruana Cayetano Heredia Research Ethics Committee (Lima, Peru). Participants provided written, informed consent.

Submitted March 16, 2011; accepted November 18, 2011. Prepublished online as *Blood* First Edition paper, November 30, 2011; DOI: 10.1182/blood-2011-03-341776.

*N.P.T., S.L., and T.G.S. contributed equally to this work.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

Table 1. Baseline characteristics of study participants

	Iron group (n = 12)	Saline group (n = 12)
Age, y	32 ± 11	33 ± 9
Height, cm	166 ± 4	166 ± 6
Weight, kg	73 ± 13	72 ± 9
End-tidal PO ₂ , mmHg	104 ± 8	106 ± 4
Arterial oxygen saturation, %	99 ± 1	99 ± 1
Hematocrit, %	47 ± 2	45 ± 2
Ferritin, ng/mL	96 ± 69	90 ± 34
Transferrin saturation, %	32 ± 9	31 ± 14
Hepcidin, ng/mL	14 ± 11	14 ± 11
Growth differentiation factor-15, pg/mL	532 ± 230	515 ± 177
Erythropoietin, mU/mL	4.4 ± 1.9	7.0 ± 4.2
Soluble transferrin receptor, nmol/L	20 ± 5	17 ± 4

Values are mean ± SD.
PO₂ indicates partial pressure of oxygen.

Iron loading before ascent to high altitude

Twenty-four healthy male volunteers who lived at sea level (age 32 ± 10 years, mean ± SD) were recruited in Lima, Peru, where baseline (0 hours) blood tests were performed. On the morning of day 1, volunteers were randomized to receive an intravenous infusion of either iron(III) sucrose (200 mg in 100 mL over the course of 30 minutes; Vifor Inc) or saline placebo (100 mL over the course of 30 minutes). Within 2 hours of infusion, volunteers began the 8-hour ascent by road to Cerro de Pasco (4340 m). Blood tests were repeated on arrival, and on the mornings of days 2, 3, and 7.

Hematologic indices of iron homeostasis

At each time point, hematocrit was assessed by microcentrifugation. Serum for measurement of transferrin saturation and ferritin concentration was stored at 4°C and analyzed within 72 hours. Serum was stored at -20°C for erythropoietin ELISA (Medlab, Lima, Peru). Plasma was stored at -20°C for hepcidin ELISA (Bachem) and sTfR, GDF15, and IL-6 ELISA (R&D Systems). Hepcidin and IL-6 values for one volunteer at one time point




Figure 1. Changes in arterial oxygen saturation (arterial SpO₂, measured by pulse oximetry), hematologic markers of iron homeostasis, serum EPO, plasma sTfR, and GDF15 after a rapid ascent to high altitude. Baseline (0 hours) measurements were made in Lima, Peru (133 m above sea level), after which volunteers received an intravenous infusion of either 200 mg of iron sucrose (closed symbols, n = 12) or saline (open symbols, n = 12) and then ascended rapidly to high altitude (4340 m). *Significant difference between the change from baseline in the iron group, compared with the saline group (unpaired Student *t* test, *P* < .01). †, ‡Significant change from baseline within the iron or saline group, respectively (paired Student *t* test, *P* < .05). Note that initial apparent "oversaturation" of transferrin (calculated transferrin saturation > 100%) can occur with intravenous administration of iron, reflecting the transient excess of serum iron. Symbols represent mean ± SEM.

(day7) were excluded from analysis because this volunteer developed a severe upper respiratory tract infection, which is known to enhance hepcidin expression.^{17,18}

Statistical analysis

Contrasts between iron infusion and placebo were drawn by the use of repeated-measures ANOVA, unless otherwise indicated. Specific time points within and between groups were compared with the use of paired and unpaired Student *t* tests, respectively. Differences were considered significant if $P < .05$. Values are presented as mean \pm SEM.

Results and discussion

Baseline characteristics of volunteers in the iron and saline groups are summarized in Table 1. The groups were well-matched and all participants were iron-replete at baseline. Figure 1 shows that in volunteers given a placebo infusion before ascent, hepcidin decreased by the first morning at altitude ($P < .01$). This decrease occurred substantially (> 24 hours) before any change in transferrin saturation, serum ferritin concentration, or sTfR concentration, suggesting that it did not require the development of iron deficiency. The decrease in hepcidin was temporally associated with significant increases in EPO and GDF15 levels ($P < .01$ in each case), but further studies will be required to establish whether the association was causative in either case. Notably, there was no significant correlation between changes in hepcidin and those of GDF15 or EPO at any time point, and plasma levels of GDF15 were approximately 100-fold lower than those of patients with β -thalassemia, and 10-fold lower than those shown to suppress hepcidin in vitro.¹² Plasma sTfR was stable for the first 48 hours at altitude, but elevated by day 7 ($P < .01$). This result is in keeping with the observation that sTfR is primarily an index of erythropoietic activity in iron-replete subjects.^{19,20}

Intravenous iron loading induced a rapid, but transient, increase in transferrin saturation together with a more gradual, but persistent, increase in serum ferritin. The variation in hepcidin over time appeared very similar to that for transferrin saturation, but it was dissociated from the variation in ferritin over time. Of note, plasma hepcidin concentration in the group that received iron was markedly suppressed by day 7, compared with baseline ($P < .01$), despite a mean ferritin > 200 ng/mL. Iron loading had no signifi-

cant effect on EPO, sTfR, or GDF15 levels at altitude, compared with saline ($P > .2$).

Inflammation can induce hepcidin expression via IL-6 production.^{18,21} In the current study, plasma IL-6 was above the normal range for only 1 of 119 samples analyzed. The IL-6 concentration was below the lowest assay standard value in 79% of samples from volunteers receiving iron, and 69% of samples from those receiving saline ($P > .2$, χ^2 test).

Taken together, these findings demonstrate that the suppression of hepcidin by hypoxia is too fast to be generated by a reduction in iron availability, suggesting that erythropoiesis and/or hypoxia per se are the major physiologic regulators in this setting. Iron loading can overcome the suppression of hepcidin by hypoxia during the period for which transferrin saturation remains elevated, but not during the entire period for which ferritin remains elevated. This finding is consistent with the notion that transferrin saturation is the major signal linking iron availability to hepcidin up-regulation,^{5,8,9} but intrahepatic iron has also been reported to enhance hepcidin expression directly.^{22,23} The latter phenomenon could explain the slightly higher hepcidin levels after 48 hours in the iron-loaded group, compared with the saline group, despite similar transferrin saturations.

Acknowledgments

The authors thank the volunteers who took part in the study.

This research was supported by the Wellcome Trust and the Higher Education Funding Council for England.

Authorship

Contribution: N.P.T., S.L., T.G.S., F.L.-V., K.L.D., D.R.M., and P.A.R. designed the research; N.P.T., S.L., T.G.S., C.P., A.H.N., and M.R.-C. performed experiments; and N.P.T., S.L., T.G.S., D.R.M., and P.A.R. analyzed and interpreted the data and wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Professor Peter Robbins, Department of Physiology, Anatomy & Genetics, University of Oxford, Oxford OX1 3PT, United Kingdom; e-mail: peter.robbins@dpag.ox.ac.uk.

References

- Bothwell TH, Pirzio-Biroli G, Finch CA. Iron absorption. I. Factors influencing absorption. *J Lab Clin Med*. 1958;51(1):24-36.
- Finch C. Regulators of iron balance in humans. *Blood*. 1994;84(6):1697-1702.
- Nicolas G, Bennoun M, Devaux I, et al. Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice. *Proc Natl Acad Sci U S A*. 2001; 98(15):8780-8785.
- Nemeth E, Tuttle MS, Powelson J, et al. Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science*. 2004;306(5704):2090-2093.
- Lin L, Valore EV, Nemeth E, Goodnough JB, Gabayan V, Ganz T. Iron transferrin regulates hepcidin synthesis in primary hepatocyte culture through hemojuvelin and BMP2/4. *Blood*. 2007; 110(6):2182-2189.
- Girelli D, Trombini P, Busti F, et al. A time course of hepcidin response to iron challenge in patients with HFE and TFR2 hemochromatosis. *Haematologica*. 2011;96(4):500-506.
- Gao J, Chen J, Kramer M, Tsukamoto H, Zhang AS, Enns CA. Interaction of the hereditary hemochromatosis protein HFE with transferrin receptor 2 is required for transferrin-induced hepcidin expression. *Cell Metab*. 2009;9(3): 217-227.
- Giannetti AM, Bjorkman PJ. HFE and transferrin directly compete for transferrin receptor in solution and at the cell surface. *J Biol Chem*. 2004; 279(24):25866-25875.
- Schmidt PJ, Toran PT, Giannetti AM, Bjorkman PJ, Andrews NC. The transferrin receptor modulates Hfe-dependent regulation of hepcidin expression. *Cell Metab*. 2008;7(3): 205-214.
- Babitt JL, Huang FW, Wrighting DM, et al. Bone morphogenetic protein signaling by hemojuvelin regulates hepcidin expression. *Nat Genet*. 2006; 38(5):531-539.
- Hentze MW, Muckenthaler MU, Andrews NC. Balancing acts: molecular control of mammalian iron metabolism. *Cell*. 2004;117(3):285-297.
- Tanno T, Bhanu NV, Oneal PA, et al. High levels of GDF15 in thalassemia suppress expression of the iron regulatory protein hepcidin. *Nat Med*. 2007;13(9):1096-1101.
- Peyssonaux C, Zinkernagel AS, Schuepbach RA, et al. Regulation of iron homeostasis by the hypoxia-inducible transcription factors (HIFs). *J Clin Invest*. 2007;117(7):1926-1932.
- Volke M, Gale DP, Maegdefrau U, et al. Evidence for a lack of a direct transcriptional suppression of the iron regulatory peptide hepcidin by hypoxia-inducible factors. *PLoS One*. 2009; 4(11):e7875.
- Lakhal S, Schödel J, Townsend AR, Pugh CW, Ratcliffe PJ, Mole DR. Regulation of type II transmembrane serine proteinase TMPRSS6 by hypoxia-inducible factors: new link between hypoxia signaling and iron homeostasis. *J Biol Chem*. 2011;286(6):4090-4097.
- Silvestri L, Pagani A, Camaschella C. Furin-mediated release of soluble hemojuvelin: a new link between hypoxia and iron homeostasis. *Blood*. 2008;111(2):924-931.

17. Piperno A, Galimberti S, Mariani R, et al. Modulation of hepcidin production during hypoxia-induced erythropoiesis in humans in vivo: data from the HIGHCARE project. *Blood*. 2011;117(10):2953-2959.
18. Nicolas G, Chauvet C, Viatte L, et al. The gene encoding the iron regulatory peptide hepcidin is regulated by anemia, hypoxia, and inflammation. *J Clin Invest*. 2002;110(7):1037-1044.
19. Townsend A, Drakesmith H. Role of HFE in iron metabolism, hereditary haemochromatosis, anaemia of chronic disease, and secondary iron overload. *Lancet*. 2002;359(9308):786-790.
20. Beguin Y. Soluble transferrin receptor for the evaluation of erythropoiesis and iron status. *Clin Chim Acta*. 2003;329(1-2):9-22.
21. Nemeth E, Rivera S, Gabayan V, et al. IL-6 mediates hypoferrremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest*. 2004;113(9):1271-1276.
22. Meynard D, Kautz L, Darnaud V, Canonne-Hergaux F, Coppin H, Roth MP. Lack of the bone morphogenetic protein BMP6 induces massive iron overload. *Nat Genet*. 2009;41(4):478-481.
23. Kautz L, Meynard D, Monnier A, et al. Iron regulates phosphorylation of Smad1/5/8 and gene expression of Bmp6, Smad7, Id1, and Atoh8 in the mouse liver. *Blood*. 2008;112(4):1503-1509.